# GENERALIZATION OF A DENSITY THEOREM OF KHINCHIN AND DIOPHANTINE APPROXIMATION

### J. BECK AND W.W.L. CHEN

ABSTRACT. The continuous version of a fundamental result of Khinchin says that a half-infinite torus line in the unit square  $[0,1]^2$  exhibits superdensity, which is a best form of time-quantitative density, if and only if the slope of the geodesic is a badly approximable number. We extend this result of Khinchin to the case when the unit torus  $[0,1]^2$  is replaced by a finite polysquare translation surface, or square tiled surface. In particular, we show that it is possible to study this very number-theoretic problem by restricting to traditional tools in number theory, using only continued fractions and the famous 3-distance theorem in diophantine approximation combined with an iterative process.

We improve on an earlier result of the authors and Yang [3] where it is shown that badly approximable numbers that satisfy a quite severe technical restriction on the digits of their continued fractions lead to superdense geodesics. Here we overcome this technical impediment.

This paper is self-contained, and the reader does not need any knowledge of dynamical systems.

#### 1. Introduction

It is well known that the distribution of the irrational rotation sequence  $n\alpha \mod 1$ ,  $n=1,2,3,\ldots$ , is intimately related to the distribution of half-infinite torus lines of slope  $\alpha$  in the unit square  $[0,1]^2$ , *i.e.*, geodesics of slope  $\alpha$  on the unit torus  $[0,1]^2$ . An old result of Khinchin [8, Theorem 26] implies the following result concerning superdensity of geodesics; an alternative proof can be found in [3, Lemma 6.1.1].

**Theorem** (Khinchin). Any half-infinite geodesic is superdense on the unit torus  $[0,1]^2$  if and only if the slope of the geodesic is a badly approximable number.

Recall that an irrational number  $\alpha$ , with continued fraction

$$\alpha = [a_0; a_1, a_2, a_3, \ldots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_2 + \cdots}}},$$
 (1.1)

is said to be badly approximable if there exists a constant A such that the continued fraction digits  $a_i \leq A$  for every  $i = 0, 1, 2, 3, \dots$ 

Superdensity is a time-quantitative criterion. A half-infinite geodesic  $\mathcal{L}(t)$ ,  $t \geq 0$ , equipped with the usual arc-length parametrization, is superdense on the unit torus  $[0,1]^2$  if there exists an absolute constant  $C_1 = C_1(\mathcal{L}) > 0$  such that, for every integer  $n \geq 1$ , the initial segment  $\mathcal{L}(t)$ ,  $0 \leq t \leq C_1 n$ , of the geodesic gets (1/n)-close to every point of  $[0,1]^2$ . Note that for convenience, we use the same letter  $\mathcal{L}$  to denote both the map representing the geodesic and the initial segment of the geodesic. However, the ranges of t given remove any ambiguity.

This concept of superdensity is a best possible form of time-quantitative density, in the sense that the linear length  $C_1n$  cannot be replaced by any sublinear length o(n) as  $n \to \infty$ . For a simple proof of this; see [3, Section 6.1].

A very natural number-theoretic question concerns possible extension of the result of Khinchin by replacing the unit torus  $[0,1]^2$  by a finite surface of a certain kind.

A finite polysquare region, or a finite square tiled region in the terminology of dynamical systems, is an arbitrary connected, but not necessarily simply-connected, polygon P on the plane which is tiled with unit squares, assumed to be closed, that we call the atomic squares of P, and which satisfies the following conditions:

- (i) Any two atomic squares in P either are disjoint, or intersect at a single point, or have a common edge.
- (ii) Any two atomic squares in P are joined by a chain of atomic squares where any two neighbors in the chain have a common edge.

Note that P may have *holes*, and we also allow *whole barriers* which are horizontal or vertical *walls* that consist of one or more boundary edges of atomic squares.

Furthermore, a finite polysquare region can be converted to a finite polysquare translation surface, or square tiled surface, by pairwise identification of the boundary horizontal edges and pairwise identification of the boundary vertical edges. Geodesic flow on this surface is thus 1-direction linear flow. In Figure 1.1, we show examples of polysquare translation surfaces where identified pairs of edges can be obtained from each other by perpendicular translation. The surface in the picture on the left has 5 atomic squares, whereas the surface in the picture on the right has 32 atomic squares, 2 holes as well as 2 horizontal walls and 4 vertical walls. Some 1-direction geodesic segments are shown. These depend on the detailed edge identification which we have not shown.




Figure 1.1: some finite polysquare translation surfaces

The concept of superdensity can be extended to finite polysquare surfaces in a natural way. A half-infinite geodesic  $\mathcal{L}(t)$ ,  $t \geq 0$ , equipped with the usual arc-length parametrization, is superdense on a finite polysquare surface  $\mathcal{P}$  if there exists an absolute constant  $C_1 = C_1(\mathcal{P}; \mathcal{L}) > 0$  such that, for every integer  $n \geq 1$ , the initial segment  $\mathcal{L}(t)$ ,  $0 \leq t \leq C_1 n$ , of the geodesic gets (1/n)-close to every point of  $\mathcal{P}$ .

Using traditional tools in number theory based on diophantine approximation and continued fractions, we give a proof of the following extension of the result of Khinchin.

**Theorem 1.** Let  $\mathcal{P}$  be an arbitrary finite polysquare translation surface. Then a half-infinite geodesic that does not hit a vertex of  $\mathcal{P}$  is superdense on  $\mathcal{P}$  if and only if the slope of the geodesic is a badly approximable number.

Theorem 1 is an *if and only if* type result, where one of the two implications is a straightforward corollary of Khinchin's theorem. Indeed, 1-direction geodesic flow on a finite polysquare translation surface *modulo one* becomes 1-direction geodesic flow on the unit torus  $[0,1]^2$ , which implies that a superdense geodesic must have a badly approximable slope. The hard task is to prove the converse, that every badly approximable slope leads to superdensity.

A finite polysquare translation surface may have singularities. These then make the system non-integrable and the analysis much harder. A pioneering result in this direction concerns geodesics on a large class of surfaces, which we state below in the special case of finite polysquare translation surfaces. This result is time-qualitative in nature, in that it does not give any indication on how long it takes for the geodesic to get within a given distance of a given point in  $\mathcal{P}$ . Indeed, the traditional approach from the viewpoint of dynamical systems is based on application of results such as Birkhoff's ergodic theorem which are essentially time-qualitative in nature. Lacking an error term, they do not appear to lead naturally to time-quantitative statements.

**Theorem** (Katok–Zemlyakov [6]). Any half-infinite geodesic on a finite polysquare translation surface  $\mathcal{P}$  with irrational slope is dense unless it hits a vertex of  $\mathcal{P}$  and becomes undefined.

In a recent series of papers [1, 2, 3, 4], the authors and their co-authors are able to establish many results concerning the long-term time-quantitative behavior in many flat systems concerning 1-direction geodesic flow on surfaces where the faces have zero curvature. In particular, a weaker form of Theorem 1 is established, where it is shown that superdensity follows if the slope  $\alpha$  given by (1.1) satisfies the additional technical requirement that the digits  $a_0, a_1, a_2, a_3, \ldots$  are all integer multiples of the street-LCM of the finite polysquare translation surface under consideration. The street-LCM of a finite polysquare translation surface is the lowest common multiple of the lengths of the horizontal and the vertical streets of the surface. While this excludes many badly approximable slopes, the method nevertheless gives an uncountable set of slopes which guarantee superdensity. In Theorem 1, we remove this technical impediment.

The proof of Theorem 1 here, however, is completely different from our earlier technique. However, the two different approaches share a common characteristic, in that neither is based on the traditional application of ergodic theory in the earlier study of density and uniformity using traditional techniques in dynamical systems. Instead, we appeal to a non-ergodic approach, and base our arguments on number theory and combinatorics.

We thus have two methods to prove superdensity. They are not comparable, and have different advantages. For instance, the shortline method developed in [3] works beyond geodesics on polysquare translation surfaces, and can give superdensity for geodesics on any regular polygon surface. We do not see how we can establish such a result with the method of this paper.

Our proof here of Theorem 1 is elementary but not simple. We therefore start by illustrating the ideas by studying the special case of the L-surface, which is arguably the simplest non-integrable polysquare translation surface.

The picture on the left in Figure 1.2 shows the L-shape region composed of 3 unit squares. Furthermore, it shows the L-graph, which is an undirected planar graph with 8 vertices, 5 horizontal edges and 5 vertical edges. Using this picture and identifying the edges, we can reduce the number of vertices. On identifying the edges  $h_1$ , we see that A = G and B = F. On identifying the edges  $h_2$ , we see that

B = E and C = D. On identifying the edges  $v_1$ , we see that A = C and H = D. On identifying the edges  $v_2$ , we see that H = E and G = F. Thus all the vertices are identified with each other, and we have essentially only 1 vertex. This single vertex is a split singularity point of the geodesic flow, explaining why it is a *non-integrable* system; see the two geodesics in the picture on the right in Figure 1.2.

Furthermore, the surface has 3 faces and, after identification, 6 edges, so the Euler characteristic is  $\chi = 1 - 6 + 3 = -2$ . The genus g is obtained from the well known formula  $g = 1 - (\chi/2) = 1 + 1 = 2$ , and so the surface is homeomorphic to a 2-holed torus using the classification theorem of closed surfaces. We call this the *L-surface*. It is equipped with a flat metric, and the curvature is zero on every square face. The two geodesics in the picture on the right in Figure 1.2 illustrate why the vertex E, and hence every other vertex, is a split singularity of the geodesic flow. The L-surface is classified as a Riemann surface with a singular point.




Figure 1.2: the L-surface and two geodesics

As an analog of half-infinite geodesics on the unit torus  $[0,1]^2$ , we can consider half-infinite geodesics on the L-surface. A particle moves on the geodesic with unit speed, so that time equals distance. If a geodesic on the L-surface has irrational slope and never hits the singular point, then in the L-shape it is represented as a union of infinitely many *parallel* line segments. A geodesic is uniquely determined by one of its points and its constant velocity vector, just like a geodesic on the unit torus  $[0,1]^2$ .

We shall first prove the following special case of Theorem 1.

**Theorem 2.** If a half-infinite geodesic that does not hit a vertex of the L-surface has a slope that is a badly approximable number, then it is superdense on the L-surface.

We can summarize the proof of Theorem 2, and hence also Theorem 1, in a nutshell as a *careful* use of the classical tool of continued fractions.

Before we begin our proof of Theorem 2, however, we make some comments.

First of all, the situation is completely different if we consider uniform distribution instead of density.

Corresponding to superdensity, we can define superuniformity as a best form of time-quantitative uniformity, where the relevant discrepancy is of logarithmic size compared to the length of the geodesic. For half-infinite geodesics on the unit torus  $[0,1]^2$ , it follows from time-quantitative extensions of the famous Kronecker–Weyl equidistribution theorem that a geodesic on the unit torus is superuniform if the slope is a badly approximable number. Actually, superuniformity is equivalent to the condition that the Cesaro-mean of the continued fraction digits of the slope of the geodesic is bounded.

The situation is very different if we consider instead half-infinite geodesics on the L-surface. While any half-infinite geodesic with a quadratic irrational slope is uniformly distributed on the L-surface, the rate of convergence to uniformity can be vastly different for two distinct quadratic irrational slopes. For example, it is shown in [1, 2] that a half-infinite geodesic of slope  $\alpha = \sqrt{2}$  on the L-surface is superuniform, whereas a half-infinite geodesic of slope  $\alpha = 1 + \sqrt{2}$  on the L-surface exhibits discrepancy greater than random square-root size.

Secondly, Theorem 1 has analogs for billiard orbits in finite polysquare regions and for geodesics on surfaces of finite simply-connected polycube regions. Billiard orbits in finite polysquare regions and geodesics on finite polysquare surfaces are related by the concept of unfolding due to König and Szücs [9], first introduced to the unit square, leading to 4-fold covering by reflection across a horizontal axis and across a vertical axis. For an illustration, see also [1, Section 1.3]. As to surfaces of simply-connected polycube regions, the simplest example is the surface of the unit cube. Geodesic flow on such a surface is 4-direction geodesic flow, and it can be related to 1-direction geodesic flow on a surface obtained by combining 4 rotated copies of this surface in a suitable way. For an illustration, see [4, Example 7.2.4].

Finally, it has been drawn to our attention that there is perhaps a possibility of establishing results such as Theorem 1 by using the ideas of Teichmüller dynamics, a very powerful tool in dynamical systems. Such techniques are beyond the reach of many who are not experts in that area. Our primary aim in this paper is to show that traditional number-theoretic techniques, involving only continued fractions and diophantine approximation combined with an iterative process, are sufficient for our needs. Nevertheless we make some brief comments at the end of this paper on work using this alternative approach.

## 2. Some prerequisites

Without loss of generality, we assume that the slope of the geodesic is greater than 1. Suppose that the geodesic has slope  $1/\alpha$ , where  $0 < \alpha < 1$  is irrational.

Our first tool is an interval exchange transformation  $T = T_{\alpha}$  which encodes the information concerning the particular order with which a geodesic of slope  $1/\alpha$  keeps hitting the horizontal edges  $h_1, h_2, h_3$ .

The first step involves identifying the horizontal edges  $h_1, h_2, h_3$  with unit intervals by making use of the correspondences

$$h_1 = [0, 1), \quad h_2 = [1, 2), \quad h_3 = [2, 3),$$

perhaps somewhat abusing notation, as shown in Figure 2.1.




Figure 2.1: representing horizontal edges of the L-surface by intervals. We now consider the piecewise linear map  $T=T_{\alpha}$  defined according to Figure 2.2.




Figure 2.2: the interval exchange transformation  $T = T_{\alpha}$ 

This is called the interval exchange transformation. More precisely, we have

$$T([0, 1 - \alpha)) = [2 + \alpha, 3), \quad T([1 - \alpha, 1)) = [1, 1 + \alpha),$$
 (2.1)

$$T([1, 2 - \alpha)) = [1 + \alpha, 2), \quad T([2 - \alpha, 2)) = [2, 2 + \alpha),$$
 (2.2)

$$T([2, 3 - \alpha)) = [\alpha, 1), \quad T([3 - \alpha, 3)) = [0, \alpha),$$
 (2.3)

where, for instance,  $T([0, 1 - \alpha)) = [2 + \alpha, 3)$  describes the  $1/\alpha$  flow mapping the part  $[0, 1 - \alpha)$  in  $h_1 = [0, 1)$  to the part  $[2 + \alpha, 3)$  in  $h_3 = [2, 3)$  linearly in the form

$$Tx = 2 + \alpha + x, \quad x \in [0, 1 - \alpha),$$

and similarly for the rest in (2.1)–(2.3).

The novelty is that T acts on the longer interval [0,3) instead of the unit interval [0,1), but if we consider T modulo 1, then it acts simply as an  $\alpha$ -shift, or irrational rotation, in the unit interval.

We next consider our main idea, which involves continued fractions. Consider an irrational number

$$\alpha = [a_0; a_1, a_2, a_3, \ldots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_2 + \cdots}}},$$
 (2.4)

where  $a_0 \ge 0$  and  $a_i \ge 1$ ,  $i = 1, 2, 3, \ldots$ , are integers. The rational numbers

$$\frac{p_k}{q_k} = \frac{p_k(\alpha)}{q_k(\alpha)} = [a_0; a_1, \dots, a_k], \quad k = 0, 1, 2, 3, \dots,$$
(2.5)

are the k-convergents of  $\alpha$ . It is well known that they give rise to the best rational approximations of the irrational number  $\alpha$ , and we have

$$\frac{p_0}{q_0} < \frac{p_2}{q_2} < \frac{p_4}{q_4} < \dots < \alpha < \dots < \frac{p_5}{q_5} < \frac{p_3}{q_3} < \frac{p_1}{q_1}. \tag{2.6}$$

Let ||y|| denote the distance of a real number y from the nearest integer. We shall make use of the fact that for an irrational number  $\alpha$  the sequence

$$\min_{1 \le k \le n} ||k\alpha||, \quad n = 1, 2, 3, \dots,$$

is well described by the continued fraction expansion of  $\alpha$ .

For every k = 0, 1, 2, 3, ..., we have

$$||q\alpha|| \geqslant ||q_k\alpha||, \quad 1 \leqslant q < q_{k+1},$$

$$||q_{k+1}\alpha|| < ||q_k\alpha||,$$

$$(2.7)$$

as well as

$$\frac{1}{q_{k+1} + q_k} \leqslant ||q_k \alpha|| \leqslant \frac{1}{q_{k+1}}.$$
(2.8)

Indeed, the sequences  $p_k$  and  $q_k$ ,  $k = 0, 1, 2, 3, \ldots$ , are given by the initial values

$$p_0 = a_0, \quad p_1 = a_1 a_0 + 1, \quad q_0 = 1, \quad q_1 = a_1,$$

and the recurrence relations

$$p_{k+1} = a_{k+1}p_k + p_{k-1}, \quad q_{k+1} = a_{k+1}q_k + q_{k-1}, \quad k \geqslant 1.$$
 (2.9)

We also have

$$p_{k-1}q_k - q_{k-1}p_k = (-1)^k, \quad k \geqslant 1.$$

On the other hand, using (2.6) and (2.9), it is easy to show that

$$||q_{k+1}\alpha|| + a_{k+1}||q_k\alpha|| = ||q_{k-1}\alpha||. \tag{2.10}$$

We need the following result.

**Theorem** (3-distance theorem). Consider the n + 1 numbers  $0, \alpha, 2\alpha, 3\alpha, \ldots, n\alpha$  modulo 1 in the unit torus/circle [0,1), leading to an (n+1)-partition. This partition exhibits at most 3 different distances between consecutive points. Furthermore, every positive integer n can be expressed uniquely in the form

$$n = \mu q_k + q_{k-1} + r$$
, with  $1 \leqslant \mu \leqslant a_{k+1}$  and  $0 \leqslant r < q_k$ ,

in terms of the continued fraction (2.4) of  $\alpha$  and its convergents (2.5), with the convention that  $q_{-1} = 0$ . Then

- (i) the distance  $||q_k\alpha||$  shows up precisely  $n+1-q_k$  times;
- (ii) the distance  $||q_{k-1}\alpha|| \mu ||q_k\alpha||$  shows up precisely r+1 times; and
- (iii) the distance  $||q_{k-1}\alpha|| (\mu 1)||q_k\alpha||$  shows up precisely  $q_k r 1$  times.

This surprising geometric fact, formulated as a conjecture by Steinhaus, has many proofs, by Sós [13, 14], Swierczkowski [17], Surányi [16], Halton [5] and Slater [12], with others published more recently.

# 3. Proof of Theorem 2

Given an integer  $k \ge 1$ , let  $\mathcal{A}_k(\alpha)$  denote the partition of the unit torus/circle [0,1) with  $q_{k+1} = q_{k+1}(\alpha)$  division points  $\{q\alpha\}, -1 \le q \le q_{k+1} - 2$ , where  $\{x\}$  denotes the fractional part of a real number x. Note that the choices q = -1, 0 in  $\{q\alpha\}$  represent the dangerous endpoints of the special intervals  $[0, 1 - \alpha)$  and  $[1 - \alpha, 1)$  in (2.1). These are the two singularities of the interval exchange transformation T restricted to the interval  $0 \le x < 1$ , in the sense that both 0 and  $T(1 - \alpha) = 1$  represent the split singularity of the L-surface.

A consequence of the special choice  $n = q_{k+1} - 1$  is that the 3-distance theorem simplifies to a 2-distance theorem. This in turn leads to some very useful information concerning the distances between the consecutive points of the  $q_{k+1}$ -partition  $\mathcal{A}_k(\alpha)$  of the unit torus/circle [0, 1). Indeed, using the second recurrence relation in (2.9), we have

$$n = q_{k+1} - 1 = a_{k+1}q_k + q_{k-1} - 1 = \mu q_k + q_{k-1} + r,$$

with  $\mu = a_{k+1} - 1$  and  $r = q_k - 1$ . Since  $q_k - r - 1 = 0$ , it follows from the 3-distance theorem that there are only two distances

$$||q_k \alpha||$$
 and  $||q_{k-1} \alpha|| - (a_{k+1} - 1)||q_k \alpha|| = ||q_{k+1} \alpha|| + ||q_k \alpha||,$  (3.1)

in view of (2.10).

It follows immediately from (2.7) that one of the neighbors of 0 in the partition  $\mathcal{A}_k(\alpha)$  is  $\{q_k\alpha\}$  which clearly has distance  $\|q_k\alpha\|$  from 0 in the unit torus/circle. Since  $\alpha$  is irrational, the other neighbor of 0 in the partition  $\mathcal{A}_k(\alpha)$  must have

distance  $||q_{k+1}\alpha|| + ||q_k\alpha||$  from 0 in the unit torus/circle. Simple calculation then shows that it is  $\{((a_{k+1}-1)q_k+q_{k-1})\alpha\}$ . Thus the two neighbors

$$\{q_k \alpha\}$$
 and  $\{((a_{k+1} - 1)q_k + q_{k-1})\alpha\}$ 

of 0 in the partition  $\mathcal{A}_k(\alpha)$  exhibit the two gaps in (3.1) in some order. Similarly, the two neighbors

$$\{(q_k-1)\alpha\}$$
 and  $\{((a_{k+1}-1)q_k+q_{k-1}-1)\alpha\}$ 

of  $1 - \alpha = \{-\alpha\}$  in the partition  $\mathcal{A}_k(\alpha)$  exhibit the same two gaps in (3.1) in the same order.

The union of the left and right neighborhoods of 0 in the partition  $\mathcal{A}_k(\alpha)$  has the form

$$B(0) = (-d^*, d^{**}), (3.2)$$

and the union of the left and right neighborhoods of  $1 - \alpha = \{-\alpha\}$  in the partition  $\mathcal{A}_k(\alpha)$  has a similar form

$$B(-1) = (1 - \alpha - d^*, 1 - \alpha + d^{**}), \tag{3.3}$$

due to the same order, where

$$\{d^*, d^{**}\} = \{\|q_k \alpha\|, \|q_{k+1} \alpha\| + \|q_k \alpha\|\}, \tag{3.4}$$

but we have not specified which one is which. We refer to B(0) and B(-1) as the buffer zones of the singularities 0 and  $1 - \alpha$  respectively in the partition  $\mathcal{A}_k(\alpha)$ .

We consider the special intervals

$$J_k(q) = J(\alpha; k; q) = (\{q\alpha\} - d^{**}, \{q\alpha\} + d^*), \quad 1 \le q \le q_{k+1} - 2.$$
(3.5)

Remark. These short special intervals (3.5) have three crucial properties:

- (i) They completely cover the two long intervals  $(0, 1 \alpha)$  and  $(1 \alpha, 1)$ .
- (ii) They avoid the singularities 0, 1 and  $1 \alpha$ , in view of (3.2)–(3.4).
- (iii) Any two intervals in  $(0, 1-\alpha)$  or in  $(1-\alpha, 1)$  arising from neighboring partition points exhibit substantial overlapping. More precisely, if  $1 \leq q', q'' \leq q_{k+1} 2$  are two integers such that  $\{q'\alpha\}$  and  $\{q''\alpha\}$  are neighboring points in the partition  $\mathcal{A}_k(\alpha)$ , so that both points are in the interval  $(0, 1-\alpha)$  or both points are in the interval  $(1-\alpha, 1)$ , then

$$length(J_k(q') \cap J_k(q'')) \geqslant \min\{d^*, d^{**}\} = ||q_k \alpha||.$$
(3.6)

Since the length of  $J_k(q)$  is  $2\|q_k\alpha\| + \|q_{k+1}\alpha\|$ , the trivial upper bound

$$2\|q_k \alpha\| + \|q_{k+1} \alpha\| < 3\|q_k \alpha\|$$

and (3.6) together justify the term substantial overlapping.

Since T acts on the interval/circle [0,3), for every interval  $J_k(q)$ ,  $1 \le q \le q_{k+1} - 2$ , given by (3.5), we define its 3-copy extension  $J_k(q;3)$  by

$$J_k(q;3) = J_k(q) \cup (1 + J_k(q)) \cup (2 + J_k(q)) \subset [0,3), \tag{3.7}$$

a union of  $J_k(q)$  with two of its translates.

After our preparation, we are now ready to study an orbit. Let  $\mathcal{L}_{\alpha}(S;t)$ ,  $t \geq 0$ , be a parametrized half-infinite geodesic with initial point S and slope  $1/\alpha$ , under the usual arc-length parametrization.

Let M be large, and consider the initial segment  $\mathcal{L}_{\alpha}(S;t)$ ,  $0 \leq t \leq M$ , of length M, which we denote by  $(\mathcal{L}_{\alpha}; M)$ . Suppose that

$$0 \le t_1 < t_2 < t_3 < \dots < t_m \le M, \tag{3.8}$$

where

$$t_{i+1} - t_i = \sqrt{1 + \alpha^2}, \quad 1 \leqslant i \leqslant m - 1,$$
 (3.9)

is the sequence of time instances t when the initial segment  $\mathcal{L}_{\alpha}(S;t)$ ,  $0 \leq t \leq M$ , intersects the union  $h_1 \cup h_2 \cup h_3 = [0,3)$  of the 3 horizontal edges of the L-surface in Figure 2.1. For notational simplicity let

$$y_i = \mathcal{L}_{\alpha}(S; t_i) \in [0, 3), \quad 1 \leqslant i \leqslant m, \tag{3.10}$$

denote these intersection points.

Using the interval exchange transformation  $T = T_{\alpha} : [0,3) \to [0,3)$ , we see that any two time-consecutive intersection points are governed by the simple relation

$$T(y_i) = y_{i+1}, \quad 1 \le i \le m-1.$$
 (3.11)

**Lemma 3.1.** Suppose that  $J_k(\ell_1)$  is a special interval of the form (3.5), and there exists  $r(\ell_1) \in \{0, 1, 2\}$  such that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap (r(\ell_1) + J_k(\ell_1)) = \emptyset, \tag{3.12}$$

where  $\{y_i: 1 \leq i \leq m\}$  is the set of intersection points defined in (3.10). Then for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ , we have

$$\{y_i: q_{k+1} \leqslant i \leqslant m - q_{k+1}\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$
 (3.13)

*Proof.* Since  $q_{k+1} \ge 1$  and  $m - q_{k+1} \le m$ , it follows trivially from (3.12) that (3.13) holds for h = 0.

Combining (3.11) and (3.12), we see that

$$\{y_i: 2 \leqslant i \leqslant m\} \cap T(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$

Iterating this argument, we see that for every integer  $1 \leq h \leq q_{k+1} - 2 - \ell_1$ , we have

$$\{y_i: 1+h \le i \le m\} \cap T^h(r(\ell_1)+J_k(\ell_1)) = \emptyset.$$

Since  $q_{k+1} \ge q_{k+1} - 1 - \ell_1 \ge h + 1$  and  $m - q_{k+1} \le m$ , it follows that (3.13) holds for every integer  $1 \le h \le q_{k+1} - 2 - \ell_1$ .

For every negative integer  $1 - \ell_1 \leq h \leq -1$ , using the inverse transformation  $T^{-1}$ , combining (3.11) and (3.12), and iterating, we have

$$\{y_i: 1 \leqslant i \leqslant m+h\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$

Since  $q_{k+1} \ge 1$  and  $m+h \ge m+1-\ell_1 \ge m+1-q_{k+1} \ge m-q_{k+1}$ , it follows that (3.13) holds for every integer  $1-\ell_1 \le h \le -1$ .

The proof of the lemma is now complete.

*Remark.* We often refer to the deduction of (3.13) from (3.12) as a T-power extension argument.

Let  $1 \leq \ell_1 \leq q_{k+1} - 2$  be fixed.

For notational convenience, for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ , we write

$$T^{h}(r(\ell_1) + J_k(\ell_1)) = r(\ell_1 + h) + J_k(\ell_1 + h).$$
(3.14)

Note that (3.14) defines r(q) for every integer  $1 \leqslant q \leqslant q_{k+1} - 2$ . Furthermore, combining (3.13) and (3.14), we have

$$\{y_i : q_{k+1} \le i \le m - q_{k+1}\} \cap (r(q) + J_k(q)) = \emptyset$$
 (3.15)

for every integer  $1 \leq q \leq q_{k+1} - 2$ .

Suppose that  $I_0 \subset [0,3)$  is  $(\mathcal{L}_{\alpha}; M)$ -free, so that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap I_0 = \emptyset,$$

where  $\{y_i : 1 \leq i \leq m\}$  is the set of intersection points defined in (3.10). Let k be an integer, and suppose that  $J_k(\ell_1)$  is a special interval of the form (3.5), and there exists  $r(\ell_1) \in \{0, 1, 2\}$  such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0. \tag{3.16}$$

Then (3.12) holds.

Remark. We shall later choose an optimal value of k for which (3.16) holds.

We distinguish a few cases according to the special relations between various sets of intersection points and various special intervals. We take advantage of the substantial overlapping of the short special intervals  $J_k(q)$  defined by (3.5).

Recall that if  $1 \leq q', q'' \leq q_{k+1} - 2$  are two integers such that  $\{q'\alpha\}$  and  $\{q''\alpha\}$  are neighboring points in the partition  $\mathcal{A}_k(\alpha)$ , so that both points are in the interval  $(0, 1 - \alpha)$  or both points are in the interval  $(1 - \alpha, 1)$ , then combining (2.8) and (3.6), we have

length
$$(J_k(q') \cap J_k(q'')) \ge ||q_k \alpha|| \ge \frac{1}{q_{k+1} + q_k} > \frac{1}{2q_{k+1}}$$
.

Recall also from (2.8), (3.4) and (3.5) that

length
$$(J_k(q)) = 2||q_k\alpha|| + ||q_{k+1}\alpha|| < 3||q_k\alpha|| \le \frac{3}{q_{k+1}},$$

so that

$$length(J_{k+8}(q)) < \frac{3}{q_{k+9}}. (3.17)$$

On the other hand, a trivial deduction from (2.9) gives

$$q_{k+2} \geqslant q_{k+1} + q_k \geqslant 2q_k,$$

so that iterating this a few times, we conclude that

$$q_{k+9} \geqslant 2q_{k+7} \geqslant 4q_{k+5} \geqslant 8q_{k+3} \geqslant 16q_{k+1}.$$
 (3.18)

Combining (3.17) and (3.18), we conclude that the intersection  $J_k(q') \cap J_k(q'')$  must contain a special interval of the type  $J_{k+8}(q)$  for some  $1 \leq q \leq q_{k+9} - 2$ . We split the argument into two complementary cases.

Case 1A. The following intersection property holds. For every

$$J_{k+8}(\ell) \subset J_k(q)$$
 and  $r \in \{0, 1, 2\},\$ 

with  $r \neq r(q)$  given by (3.14), we have

$$\{y_i : q_{k+1} \le i \le m - q_{k+1}\} \cap (r + J_{k+8}(\ell)) \ne \emptyset.$$
 (3.19)

Lemma 3.2. Case 1A is impossible.

Case 1B. There exist

$$J_{k+8}(\ell_2) \subset J_k(q^+)$$
 and  $r_1 \in \{0, 1, 2\},$ 

with  $r_1 \neq r(q^+)$  given by (3.14), such that

$$\{y_i: q_{k+1} \le i \le m - q_{k+1}\} \cap (r_1 + J_{k+8}(\ell_2)) = \emptyset.$$
 (3.20)

We now continue with this particular value of  $\ell_2$ .

Since  $J_{k+8}(\ell_2) \subset J_k(q^+)$ , it clearly follows from (3.15) that

$$\{y_i: q_{k+1} \leqslant i \leqslant m - q_{k+1}\} \cap (r(q^+) + J_{k+8}(\ell_2)) = \emptyset.$$
 (3.21)

Since (3.20) and (3.21) are analogs of (3.12), the *T*-power expansion argument in the proof of Lemma 3.1 shows that for every integer  $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$ , we have

$$\{y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}\} \cap T^h(r_1 + J_{k+8}(\ell_2)) = \emptyset, \tag{3.22}$$

$$\{y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}\} \cap T^h(r(q^+) + J_{k+8}(\ell_2)) = \emptyset.$$
 (3.23)

For notational convenience, for every integer  $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$ , we write

$$T^{h}(r(q^{+}) + J_{k+8}(\ell_{2})) = r^{\star}(\ell_{2} + h) + J_{k+8}(\ell_{2} + h), \tag{3.24}$$

$$T^{h}(r_1 + J_{k+8}(\ell_2)) = r^{\star \star}(\ell_2 + h) + J_{k+8}(\ell_2 + h). \tag{3.25}$$

Then combining (3.22)–(3.25), we have

$$\{y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}\} \cap (r^*(q) + J_{k+8}(q)) = \emptyset, \tag{3.26}$$

$$\{y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}\} \cap (r^{\star \star}(q) + J_{k+8}(q)) = \emptyset, \tag{3.27}$$

for every integer  $1 \leqslant q \leqslant q_{k+9} - 2$ . Clearly

$$r^{\star}(q) \neq r^{\star \star}(q), \quad 1 \leqslant q \leqslant q_{k+9} - 2.$$

We now split Case 1B into two complementary cases.

Case 2A. The following intersection property holds. For every

$$J_{k+16}(\ell) \subset J_{k+8}(q)$$
 and  $r \in \{0, 1, 2\},$ 

with  $r \neq r^{\star}(q), r^{\star\star}(q)$  given by (3.24) and (3.25), we have

$${y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}} \cap (r + J_{k+16}(\ell)) \neq \emptyset.$$

Lemma 3.3. Case 2A is impossible.

Case 2B. There exist

$$J_{k+16}(\ell_3) \subset J_{k+8}(q^{++})$$
 and  $r_2 \in \{0, 1, 2\},$ 

with  $r_2 \neq r^*(q^{++}), r^{**}(q^{++})$  given by (3.24) and (3.25), such that

$$\{y_i: q_{k+1} + q_{k+9} \leqslant i \leqslant m - q_{k+1} - q_{k+9}\} \cap (r_2 + J_{k+16}(\ell_3)) = \emptyset.$$
 (3.28)

Lemma 3.4. If Case 2B holds, then

$$m \leqslant 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 6. \tag{3.29}$$

Proof of Theorem 2. Suppose that  $I_0 \subset [0,3)$  is  $(\mathcal{L}_{\alpha}; M)$ -free, so that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap I_0 = \emptyset.$$

Let  $J_k(\ell_1) = J(\alpha; k; \ell_1)$  be the longest special interval of the form (3.5) such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0$$
 for some  $\ell_1$  and  $r(\ell_1) \in \{0, 1, 2\}$ .

Then

$$length(I_0) < 4(||q_{k-1}\alpha|| + ||q_k\alpha||), \tag{3.30}$$

since otherwise there exists a longer special interval

$$r(\ell) + J_{k-1}(\ell) \subset I_0$$
 for some  $\ell$  and  $r(\ell) \in \{0, 1, 2\}$ ,

a contradiction. Combining (3.30) with (2.8), we have

length(
$$I_0$$
)  $< \frac{4}{q_k} + \frac{4}{q_{k+1}} < \frac{8}{q_k}$ . (3.31)

On the other hand, it follows from (3.8) and (3.9) that

$$M \leqslant (m+1)\sqrt{1+\alpha^2}.\tag{3.32}$$

Also, in view of Lemmas 3.2–3.4, it is clear that the bound (3.29) holds. Finally, recall that  $0 < \alpha < 1$  and

$$\alpha = [a_1, a_2, a_3, \dots] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}$$

is badly approximable, so there exists a constant A such that the continued fraction digits  $a_i \leq A$  for every  $i = 1, 2, 3, \ldots$  It follows from (2.9) that

$$q_{k+1} < q_{k+9} < q_{k+17} \le (A+1)^{17} q_k.$$
 (3.33)

Combining (3.29), (3.32) and (3.33), we see that

$$M \leq (2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 7)\sqrt{1 + \alpha^2} < 7(A+1)^{17}q_k\sqrt{2}. \tag{3.34}$$

It now follows from (3.31) and (3.34) that a geodesic segment  $\mathcal{L}_{\alpha}(S;t)$ ,  $0 \leq t \leq M$ , of length  $M = 7(A+1)^{17}q_k\sqrt{2}$  must intersect every subinterval I of  $h_1 \cup h_2 \cup h_3$  with length I is a constant independent of I, this establishes superdensity of the half-infinite geodesic.

### 4. Proof of Lemmas 3.2–3.4

Before we present the proof of our main lemmas, we begin by investigating a simple situation which serves to illustrate our method.

Recall that  $r(\ell_1)$  is given by Lemma 3.2.

Simple Case. There exist integers  $r^*(\ell_1)$  and  $r^{**}(\ell_1)$  such that

- (1)  $r(\ell_1), r^*(\ell_1), r^{**}(\ell_1)$  form a permutation of 0, 1, 2;
- (2)  $\{y_i : 1 \leq i \leq m\} \cap (r^*(\ell_1) + J_k(\ell_1)) = \emptyset$ ; and
- (3)  $\{y_i : 1 \leq i \leq m\} \cap (r^{**}(\ell_1) + J_k(\ell_1)) = \emptyset.$

**Lemma 4.1.** If the Simple Case holds, then  $m \leq 2q_{k+1} + 6$ .

*Proof.* For notational simplicity, we write

$$Q(k;m) = \{ y_i : q_{k+1} \leqslant i \leqslant m - q_{k+1} \}. \tag{4.1}$$

Since the properties (2) and (3) in the Simple Case are analogs of (3.12), we can repeat the *T*-power extension argument in Lemma 3.1 and conclude that for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ , we have

$$Q(k;m) \cap T^h(r^*(\ell_1) + J_k(\ell_1)) = \emptyset, \tag{4.2}$$

$$Q(k;m) \cap T^{h}(r^{**}(\ell_{1}) + J_{k}(\ell_{1})) = \emptyset.$$
(4.3)

As in (3.14), we write

$$T^{h}(r^{*}(\ell_{1}) + J_{k}(\ell_{1})) = r^{*}(\ell_{1} + h) + J_{k}(\ell_{1} + h), \tag{4.4}$$

$$T^{h}(r^{**}(\ell_1) + J_k(\ell_1)) = r^{**}(\ell_1 + h) + J_k(\ell_1 + h), \tag{4.5}$$

for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ . Note that (4.4) and (4.5) define  $r^*(q)$  and  $r^{**}(q)$  respectively for every integer  $1 \leq q \leq q_{k+1} - 2$ . Recall next that r(q)

is defined by (3.14). Indeed, using the notation (3.14), it is easy to check that the assertion (3.13) for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$  implies that

$$Q(k;m) \cap (r(\ell_1 + h) + J_k(\ell_1 + h)) = \emptyset$$
(4.6)

for every integer  $1 - \ell_1 \leqslant h \leqslant q_{k+1} - 2 - \ell_1$ .

Combining (4.2)–(4.6) for every integer  $1 - \ell_1 \leqslant h \leqslant q_{k+1} - 2 - \ell_1$ , we deduce that for every integer  $1 \leqslant q \leqslant q_{k+1} - 2$ , we have

$$Q(k;m) \cap (r^*(q) + J_k(q)) = \emptyset, \tag{4.7}$$

$$Q(k;m) \cap (r^{**}(q) + J_k(q)) = \emptyset, \tag{4.8}$$

$$Q(k;m) \cap (r(q) + J_k(q)) = \emptyset. \tag{4.9}$$

Also, in view of the property (1) in the Simple Case, it is clear that  $r(q), r^*(q), r^{**}(q)$  form a permutation of 0, 1, 2 for every integer  $1 \leq q \leq q_{k+1} - 2$ .

By definition, the 3-copy extensions  $J_k(q;3)$ ,  $1 \leq q \leq q_{k+1} - 2$ , give rise to 6 continuous chains of overlapping intervals in the torus/circle [0,3) such that the 6 chains completely cover the 6 intervals

$$(0, 1 - \alpha), (1 - \alpha, 1), (1, 2 - \alpha), (2 - \alpha, 2), (2, 3 - \alpha), (3 - \alpha, 3),$$

and there are only 6 points in [0,3) that are not covered by the 6 chains, namely

$$0, 1-\alpha, 1, 2-\alpha, 2, 3-\alpha.$$

Combining (4.7)–(4.9) for every integer  $1 \leq q \leq q_{k+1} - 2$ , we deduce that the set Q(k;m) is not covered by the 6 chains. Indeed, if  $m \geq 2q_{k+1} + 7$ , then the set (4.1) has at least 7 distinct elements, which is more than 6, giving rise to a contradiction. We conclude therefore that, under the conditions of the Simple Case, we must have  $m \leq 2q_{k+1} + 6$ , and this completes the proof.

*Proof of Lemma 3.2.* Again, for notational simplicity, we use (4.1).

Our first step is to prove that, under the condition of Case 1A, any two neighboring 3-copy extensions  $J_k(q';3)$  and  $J_k(q'';3)$  are *synchronized* in the following precise sense: For each  $r \in \{0,1,2\}$ , we have

$$Q(k;m) \cap (r + J_k(q')) = \emptyset$$
 if and only if  $Q(k;m) \cap (r + J_k(q'')) = \emptyset$ . (4.10)

To establish this, we consider two cases.

Suppose first that r = r(q'). Using the notation (3.14), it follows from (3.13) that

$$Q(k;m) \cap (r(q') + J_k(q')) = \emptyset. \tag{4.11}$$

Assume on the contrary that

$$Q(k;m) \cap (r(q') + J_k(q'')) \neq \emptyset. \tag{4.12}$$

Then it follows from (3.15) that  $r(q'') \neq r(q')$ . On the other hand, we know that the intersection  $J_k(q') \cap J_k(q'')$  must contain a special interval of the type  $J_{k+8}(j_0)$  for some  $j_0$ , so

$$r(q') + J_{k+8}(j_0) \subset (r(q') + J_k(q')) \cap (r(q') + J_k(q'')).$$
 (4.13)

Since  $r(q') \neq r(q'')$  and  $J_{k+8}(j_0) \subset J_k(q'')$ , the condition of Case 1A is satisfied with q = q'', and (3.19) becomes

$$Q(k;m) \cap (r(q') + J_{k+8}(j_0)) \neq \emptyset.$$
 (4.14)

But (4.11) and (4.13) contradict (4.14), and so (4.12) fails. Thus our claim (4.10) holds in this case.

Suppose next that  $r \neq r(q')$ . As before, we use the fact that the intersection  $J_k(q') \cap J_k(q'')$  must contain a special interval of the type  $J_{k+8}(j_0)$  for some  $j_0$ , so

$$r + J_{k+8}(j_0) \subset (r + J_k(q')) \cap (r + J_k(q'')).$$
 (4.15)

Since  $r \neq r(q')$ , the condition of Case 1A is satisfied with q = q', and (3.19) becomes

$$Q(k;m) \cap (r + J_{k+8}(j_0)) \neq \emptyset. \tag{4.16}$$

It now follows from (4.15) and (4.16) that

$$Q(k;m) \cap (r + J_k(q')) \neq \emptyset$$
 and  $Q(k;m) \cap (r + J_k(q'')) \neq \emptyset$ ,

so that our claim (4.10) holds also in this case.

By definition, the 3-copy extensions  $J_k(q;3)$ ,  $1 \leq q \leq q_{k+1} - 2$ , give rise to 6 continuous chains of overlapping intervals in the torus/circle [0,3) such that the 6 chains completely cover the 6 intervals

$$(0,1-\alpha), (1-\alpha,1), (1,2-\alpha), (2-\alpha,2), (2,3-\alpha), (3-\alpha,3).$$
 (4.17)

The synchronization property we have just established now implies that each of the 6 long special intervals in (4.17) satisfies one of the following two properties. Either such a long special interval is *disjoint* from the set Q(k; m), or the set Q(k; m) is *dense* in such a long special interval, in the precise sense that every subinterval of length  $1/q_{k+8}$  contains a point from the set Q(k; m).

Moreover, it is not difficult to show that precisely 2 of the 6 long special intervals in (4.17) are disjoint from the set Q(k; m). To see this, choose two integers q' and q'' satisfying  $1 \leq q', q'' \leq q_{k+1} - 2$  such that

$$J_k(q') \subset (0, 1 - \alpha) \text{ and } J_k(q'') \subset (1 - \alpha, 1).$$
 (4.18)

Then it follows from (3.13) and (3.14) that

$$Q(k;m) \cap (r(q') + J_k(q')) = \emptyset$$
 and  $Q(k;m) \cap (r(q'') + J_k(q'')) = \emptyset$ . (4.19)

Now write

$$\mathcal{I}_1 = r(q') + (0, 1 - \alpha)$$
 and  $\mathcal{I}_2 = r(q'') + (1 - \alpha, 1).$  (4.20)

The synchronization property and (4.18)–(4.20) now imply that

$$Q(k;m) \cap \mathcal{I}_1 = \emptyset$$
 and  $Q(k;m) \cap \mathcal{I}_2 = \emptyset$ .

Note that the union  $\mathcal{I}_1 \cup \mathcal{I}_2$  modulo 1 is precisely the unit interval [0,1).

Now  $\mathcal{I}_1$  and  $\mathcal{I}_2$  are 2 of the 6 long special intervals in (4.17). Let  $\mathcal{I}_j$ ,  $3 \leq j \leq 6$ , denote the remaining long special intervals in (4.17). The condition of Case 1A now implies that these 4 intervals are not disjoint from Q(k;m), so that Q(k;m) is dense in each of them. Each T-image  $T(\mathcal{I}_j)$ , j=1,2, has at most 1 common point with the set Q(k;m). This is a contradiction, since the union  $T(\mathcal{I}_1) \cup T(\mathcal{I}_2)$  has a substantial intersection with the union  $\mathcal{I}_3 \cup \mathcal{I}_4 \cup \mathcal{I}_5 \cup \mathcal{I}_6$ , which implies that it must have a substantial intersection with the set Q(k;m), much more than at most 2 elements. Thus Case 1A is impossible, and this completes the proof.

*Proof of Lemma 3.3.* For notational simplicity, we write

$$Q^*(k;m) = \{ y_i : q_{k+1} + q_{k+9} \le i \le m - q_{k+1} - q_{k+9} \}. \tag{4.21}$$

We can proceed along similar lines as in the first part of the proof of Lemma 3.2 for Case 1A, and show that any two neighboring 3-copy extensions  $J_{k+8}(q';3)$  and

 $J_{k+8}(q'';3)$  are synchronized in the following precise sense: For each  $r \in \{0,1,2\}$ , we have

$$Q^*(k; m) \cap (r + J_{k+8}(q')) = \emptyset$$
 if and only if  $Q^*(k; m) \cap (r + J_{k+8}(q'')) = \emptyset$ .

By definition, the 3-copy extensions  $J_{k+8}(q;3)$ ,  $1 \le q \le q_{k+9} - 2$ , give rise to 6 continuous chains of overlapping intervals in the torus/circle [0,3) such that the 6 chains completely cover the 6 intervals (4.17). The synchronization property now implies that each of the 6 long special intervals in (4.17) satisfies one of the following two properties. Either such a long special interval is disjoint from the set  $Q^*(k;m)$ , or the set  $Q^*(k;m)$  is dense in such a long special interval, in the precise sense that every subinterval of length  $1/q_{k+16}$  contains a point from the set  $Q^*(k;m)$ .

Moreover, it is not difficult to show that precisely 4 of the 6 long special intervals in (4.17) are disjoint from the set  $Q^*(k; m)$ . To see this, choose two integers q' and q'' satisfying  $1 \leq q', q'' \leq q_{k+9} - 2$  such that

$$J_{k+8}(q') \subset (0, 1-\alpha)$$
 and  $J_{k+8}(q'') \subset (1-\alpha, 1)$ . (4.22)

Then it follows from (3.26) and (3.27) that

$$Q^*(k;m) \cap (r^*(q') + J_{k+8}(q')) = \emptyset, \quad Q^*(k;m) \cap (r^*(q'') + J_{k+8}(q'')) = \emptyset, \quad (4.23)$$

$$Q^*(k;m) \cap (r^{\star\star}(q') + J_{k+8}(q')) = \emptyset, \quad Q^*(k;m) \cap (r^{\star\star}(q'') + J_{k+8}(q'')) = \emptyset. \quad (4.24)$$

Now write

$$\mathcal{I}_1 = r^*(q') + (0, 1 - \alpha), \quad \mathcal{I}_2 = r^*(q'') + (1 - \alpha, 1),$$
 (4.25)

$$\mathcal{I}_3 = r^{\star\star}(q') + (0, 1 - \alpha), \quad \mathcal{I}_4 = r^{\star\star}(q'') + (1 - \alpha, 1).$$
 (4.26)

The synchronization property and (4.22)–(4.26) now imply that

$$Q^*(k;m) \cap \mathcal{I}_1 = \emptyset, \quad Q^*(k;m) \cap \mathcal{I}_2 = \emptyset,$$
  
 $Q^*(k;m) \cap \mathcal{I}_3 = \emptyset, \quad Q^*(k;m) \cap \mathcal{I}_4 = \emptyset.$ 

Note that the union  $\mathcal{I}_1 \cup \mathcal{I}_2 \cup \mathcal{I}_3 \cup \mathcal{I}_4$  modulo 1 is precisely the unit interval [0,1) twice.

Now  $\mathcal{I}_1, \mathcal{I}_2, \mathcal{I}_3, \mathcal{I}_4$  are 4 of the 6 long special intervals in (4.17). Let  $\mathcal{I}_j$ , j = 5, 6, denote the remaining long special intervals in (4.17). The condition of Case 2A now implies that these 2 intervals are not disjoint from  $Q^*(k;m)$ , so that  $Q^*(k;m)$  is dense in each of them. Each T-image  $T(\mathcal{I}_j)$ ,  $j = 1, \ldots, 4$ , has at most 1 common point with the set  $Q^*(k;m)$ . This is a contradiction, since the union

$$T(\mathcal{I}_1) \cup T(\mathcal{I}_2) \cup T(\mathcal{I}_3) \cup T(\mathcal{I}_4)$$

has a *substantial* intersection with the union  $\mathcal{I}_5 \cup \mathcal{I}_6$ , which implies that it must have a *substantial* intersection with the set  $Q^*(k;m)$ , much more than at most 4 elements. Thus Case 2A is impossible, and this completes the proof.

*Proof of Lemma 3.4.* For notational simplicity, we define  $Q^*(k;m)$  by (4.21), and write

$$Q^{**}(k;m) = \{ y_i : q_{k+1} + q_{k+9} + q_{k+17} \leqslant i \leqslant m - q_{k+1} - q_{k+9} - q_{k+17} \}.$$

Since (3.28) is an analog of (3.12), we can repeat the *T*-power extension argument in Lemma 3.1 and conclude that for every integer  $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$ , we have

$$Q^{**}(k;m) \cap T^h(r_2 + J_{k+16}(\ell_3)) = \emptyset.$$
(4.27)

Since  $J_{k+16}(\ell_3) \subset J_{k+8}(q^{++})$ , it follows from (3.26) and (3.27) that

$$Q^*(k;m) \cap (r^*(q^{++}) + J_{k+16}(\ell_3)) = \emptyset, \tag{4.28}$$

$$Q^*(k;m) \cap (r^{\star \star}(q^{++}) + J_{k+16}(\ell_3)) = \emptyset.$$
(4.29)

Next, note that (4.28) and (4.29) are also analogs of (3.12), so again we can repeat the *T*-power extension argument in Lemma 3.1 and conclude that for every integer  $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$ , we have

$$Q^{**}(k;m) \cap T^h(r^*(q^{++}) + J_{k+16}(\ell_3)) = \emptyset, \tag{4.30}$$

$$Q^{**}(k;m) \cap T^h(r^{**}(q^{++}) + J_{k+16}(\ell_3)) = \emptyset.$$
(4.31)

Now, for every integer  $1 - \ell_3 \leqslant h \leqslant q_{k+17} - 2 - \ell_3$ , we write

$$T^{h}(r^{\star}(q^{++}) + J_{k+16}(\ell_3)) = r^{(0)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \tag{4.32}$$

$$T^{h}(r^{\star\star}(q^{++}) + J_{k+16}(\ell_3)) = r^{(1)}(\ell_3 + h) + J_{k+16}(\ell_3 + h), \tag{4.33}$$

$$T^{h}(r_{2} + J_{k+16}(\ell_{3})) = r^{(2)}(\ell_{3} + h) + J_{k+16}(\ell_{3} + h).$$
(4.34)

Clearly it follows from the assumption of Case 2B that  $r^{(0)}(q), r^{(1)}(q), r^{(2)}(q)$  form a permutation of 0, 1, 2 for every integer  $1 \leq q \leq q_{k+17} - 2$ . Combining (4.27) and (4.30)–(4.34), we have

$$Q^{**}(k;m) \cap (r^{(0)}(q) + J_{k+16}(q)) = \emptyset, \tag{4.35}$$

$$Q^{**}(k;m) \cap (r^{(1)}(q) + J_{k+16}(q)) = \emptyset, \tag{4.36}$$

$$Q^{**}(k;m) \cap (r^{(2)}(q) + J_{k+16}(q)) = \emptyset, \tag{4.37}$$

for every integer  $1 \leqslant q \leqslant q_{k+17} - 2$ .

Note now that (4.35)–(4.37) are similar to (4.7)–(4.9) in the proof of Lemma 4.1, so we now mimic the last part of that proof.

By definition, the 3-copy extensions  $J_{k+16}(q;3)$ ,  $1 \le q \le q_{k+17} - 2$ , give rise to 6 continuous chains of overlapping intervals in the torus/circle [0,3) such that the 6 chains completely cover the 6 intervals

$$(0, 1 - \alpha), (1 - \alpha, 1), (1, 2 - \alpha), (2 - \alpha, 2), (2, 3 - \alpha), (3 - \alpha, 3),$$

and there are only 6 points in [0,3) that are not covered by the 6 chains, namely

$$0, 1-\alpha, 1, 2-\alpha, 2, 3-\alpha.$$

Combining (4.35)–(4.37) for every integer  $1 \leq q \leq q_{k+17} - 2$ , we deduce that the set  $Q^{**}(k;m)$  is not covered by the 6 chains. Indeed, if  $m \geq 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 7$ , then the set  $Q^{**}(k;m)$  has at least 7 distinct elements, which is more than 6, giving rise to a contradiction. We conclude therefore that, under the conditions of Case 2B, we must have  $m \leq 2q_{k+1} + 2q_{k+9} + 2q_{k+17} + 6$ , and this completes the proof.

### 5. Proof of Theorem 1

Consider now an arbitrary finite polysquare translation surface  $\mathcal{P}$ . We are now in a position to complete the proof of Theorem 1, and show that if the slope of a half-infinite geodesic on  $\mathcal{P}$  is a badly approximable number, then the geodesic is superdense on  $\mathcal{P}$ .

The proof is a fairly straightforward adaptation of the proof of Theorem 2, apart from the observation that the number of cases we need to consider is a function of the number of square faces of  $\mathcal{P}$ , and so can be arbitrarily large.

Without loss of generality, we assume as before that the slope of the half-infinite geodesic is greater than 1. Suppose that it has slope  $1/\alpha$ , where  $0 < \alpha < 1$  is irrational.

Our first step is to generalize the interval exchange transformation  $T = T_{\alpha}$  defined in Section 2. Suppose that the polysquare translation surface  $\mathcal{P}$  has s square faces. Each square face has a top horizontal edge and a bottom horizontal edge. Each top horizontal edge is identified with a unique bottom horizontal edge, and these give rise to s horizontal edges  $h_1, h_2, \ldots, h_s$ . We now identify these horizontal edges with unit intervals by making use of the correspondences

$$h_1 = [0, 1), \quad h_2 = [1, 2), \quad \dots, \quad h_s = [s - 1, s).$$

We now consider the piecewise linear map  $T = T_{\alpha}$  defined according to the analog of Figure 2.2 that corresponds to  $\mathcal{P}$ . More precisely, for each integer  $1 \leq j \leq s$ , there exist unique integers  $0 \leq j', j'' \leq s - 1$  such that

$$T([j-1, j-\alpha)) = [j' + \alpha, j' + 1)$$
 and  $T([j-\alpha, j)) = [j'', j'' + \alpha).$ 

Indeed, it is not difficult to see that the map  $T:[0,s)\to[0,s)$  is one-to-one and onto. While T acts on the longer interval [0,s) instead of the unit interval [0,1), if we consider T modulo 1, then it acts simply as an  $\alpha$ -shift, or irrational rotation, in the unit interval.

As before, our main idea involves continued fractions, in particular, the special case of the 3-distance theorem which becomes a 2-distance theorem. Indeed, we can repeat our discussion at the beginning of Section 3 verbatim up to the end of the paragraph preceding (3.7). In particular, for any integer  $1 \le q \le q_{k+1} - 2$ , we can define the special interval  $J_k(q)$  as in (3.5). Then analogous to (3.7), for any integer  $1 \le q \le q_{k+1} - 2$ , we define its s-copy extension  $J_k(q;s)$  by

$$J_k(q;s) = J_k(q) \cup (1 + J_k(q)) \cup \ldots \cup ((s-1) + J_k(q)) \subset [0,s),$$

a union of  $J_k(q)$  with s-1 of its translates.

Let  $\mathcal{L}_{\alpha}(t) = \mathcal{L}_{\alpha}(\mathcal{P}; S; t)$ ,  $t \ge 0$ , be a parametrized half-infinite geodesic on  $\mathcal{P}$  with initial point S and slope  $1/\alpha$ , under the usual arc-length parametrization.

Let M be large, and consider the initial segment  $\mathcal{L}_{\alpha}(t)$ ,  $0 \leq t \leq M$ , of length M, which we denote by  $(\mathcal{L}_{\alpha}; M)$ . Suppose that

$$0 \le t_1 < t_2 < t_3 < \dots < t_m \le M, \tag{5.1}$$

where

$$t_{i+1} - t_i = \sqrt{1 + \alpha^2}, \quad 1 \leqslant i \leqslant m - 1,$$
 (5.2)

is the sequence of time instances t when the initial segment  $\mathcal{L}_{\alpha}(t)$ ,  $0 \leq t \leq M$ , intersects the union  $h_1 \cup \ldots \cup h_s = [0, s)$  of the s horizontal edges of the polysquare translation surface  $\mathcal{P}$ . For notational simplicity let

$$y_i = \mathcal{L}_{\alpha}(t_i) \in [0, s), \quad 1 \leqslant i \leqslant m,$$
 (5.3)

denote these intersection points.

Using the interval exchange transformation  $T = T_{\alpha} : [0, s) \to [0, s)$ , we see that any two time-consecutive intersection points are governed by the simple relation

$$T(y_i) = y_{i+1}, \quad 1 \le i \le m-1.$$

We have the following analog of Lemma 3.1 which is easily established by the T-power extension argument.

**Lemma 5.1.** Suppose that  $J_k(\ell_1)$  is a special interval of the form (3.5), and there exists  $r(\ell_1) \in \{0, 1, ..., s-1\}$  such that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap (r(\ell_1) + J_k(\ell_1)) = \emptyset, \tag{5.4}$$

where  $\{y_i: 1 \leq i \leq m\}$  is the set of intersection points defined in (5.3). Then for every integer  $1 - \ell_1 \leq h \leq q_{k+1} - 2 - \ell_1$ , we have

$$\{y_i: q_{k+1} \le i \le m - q_{k+1}\} \cap T^h(r(\ell_1) + J_k(\ell_1)) = \emptyset.$$
 (5.5)

For notational convenience, for every integer  $1 - \ell_1 \leqslant h \leqslant q_{k+1} - 2 - \ell_1$ , we write

$$T^{h}(r(\ell_1) + J_k(\ell_1)) = r_1^{(0)}(\ell_1 + h) + J_k(\ell_1 + h).$$
(5.6)

Note that (5.6) defines r(q) for every integer  $1 \leq q \leq q_{k+1} - 2$ . Furthermore, combining (5.5) and (5.6), we have

$$\{y_i: q_{k+1} \leqslant i \leqslant m - q_{k+1}\} \cap (r_1^{(0)}(q) + J_k(q)) = \emptyset$$
(5.7)

for every integer  $1 \leqslant q \leqslant q_{k+1} - 2$ .

Suppose that  $I_0 \subset [0, s)$  is  $(\mathcal{L}_{\alpha}; M)$ -free, so that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap I_0 = \emptyset,$$

where  $\{y_i : 1 \leq i \leq m\}$  is the set of intersection points defined in (5.3). Let k be an integer, and suppose that  $J_k(\ell_1)$  is a special interval of the form (3.5), and there exists  $r(\ell_1) \in \{0, 1, \ldots, s-1\}$  such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0. \tag{5.8}$$

Then (5.4) holds. As before, we shall later choose an *optimal* value of k for which (5.8) holds.

Again, we distinguish a few cases according to the special relations between various sets of intersection points and various special intervals. We take advantage of the substantial overlapping of the short special intervals  $J_k(q)$  defined by (3.5).

Recall that if  $1 \leqslant q', q'' \leqslant q_{k+1} - 2$  are two integers such that  $\{q'\alpha\}$  and  $\{q''\alpha\}$  are neighboring points in the partition  $\mathcal{A}_k(\alpha)$ , then the intersection  $J_k(q') \cap J_k(q'')$  must contain a special interval of the type  $J_{k+8}(q)$  for some  $1 \leqslant q \leqslant q_{k+9} - 2$ . We split the argument into two complementary cases.

Write

$$Q(k; m) = \{ y_i : q_{k+1} \leqslant i \leqslant m - q_{k+1} \}.$$

Case 1A. The following intersection property holds. For every

$$J_{k+8}(\ell) \subset J_k(q)$$
 and  $r \in \{0, 1, \dots, s-1\},\$ 

with  $r \neq r_1^{(0)}(q)$  given by (5.6), we have

$$Q(k;m) \cap (r + J_{k+8}(\ell)) \neq \emptyset.$$

Case 1B. There exist

$$J_{k+8}(\ell_2) \subset J_k(q^{(1)})$$
 and  $r_1 \in \{0, 1, \dots, s-1\},\$ 

with  $r_1 \neq r_1^{(0)}(q^{(1)})$  given by (5.6), such that

$$Q(k;m) \cap (r_1 + J_{k+8}(\ell_2)) = \emptyset.$$
 (5.9)

Since  $J_{k+8}(\ell_2) \subset J_k(q^{(1)})$ , it clearly follows from (5.7) that

$$Q(k;m) \cap (r_1^{(0)}(q^{(1)}) + J_{k+8}(\ell_2)) = \emptyset.$$
(5.10)

Since (5.9) and (5.10) are analogs of (5.4), the *T*-power expansion argument shows that for every integer  $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$ , we have

$$Q^{(1)}(k;m) \cap T^h(r_1 + J_{k+8}(\ell_2)) = \emptyset, \tag{5.11}$$

$$Q^{(1)}(k;m) \cap T^{h}(r_{1}^{(0)}(q^{(1)}) + J_{k+8}(\ell_{2})) = \emptyset, \tag{5.12}$$

where, corresponding to  $Q^*(k;m)$  in Section 4, we write

$$Q^{(1)}(k;m) = \{ y_i : q_{k+1} + q_{k+9} \le i \le m - q_{k+1} - q_{k+9} \}.$$

For notational convenience, for every integer  $1 - \ell_2 \leq h \leq q_{k+9} - 2 - \ell_2$ , we write

$$T^{h}(r_{1}^{(0)}(q^{(1)}) + J_{k+8}(\ell_{2})) = r_{2}^{(0)}(\ell_{2} + h) + J_{k+8}(\ell_{2} + h), \tag{5.13}$$

$$T^{h}(r_1 + J_{k+8}(\ell_2)) = r_2^{(1)}(\ell_2 + h) + J_{k+8}(\ell_2 + h).$$
 (5.14)

Then combining (5.11)–(5.14), we have

$$Q^{(1)}(k;m) \cap (r_2^{(0)}(q) + J_{k+8}(q)) = \emptyset, \tag{5.15}$$

$$Q^{(1)}(k;m) \cap (r_2^{(1)}(q) + J_{k+8}(q)) = \emptyset, \tag{5.16}$$

for every integer  $1 \leqslant q \leqslant q_{k+9} - 2$ . Clearly

$$r_2^{(0)}(q) \neq r_2^{(1)}(q), \quad 1 \leqslant q \leqslant q_{k+9} - 2.$$

We now split Case 1B into two complementary cases.

Case 2A. The following intersection property holds. For every

$$J_{k+16}(\ell) \subset J_{k+8}(q)$$
 and  $r \in \{0, 1, \dots, s-1\},\$ 

with  $r \neq r_2^{(0)}(q), r_2^{(1)}(q)$  given by (5.13) and (5.14), we have

$$Q^{(1)}(k;m) \cap (r + J_{k+16}(\ell)) \neq \emptyset.$$

Case 2B. There exist

$$J_{k+16}(\ell_3) \subset J_{k+8}(q^{(2)})$$
 and  $r_2 \in \{0, 1, \dots, s-1\},$ 

with  $r_2 \neq r_2^{(0)}(q^{(2)}), r_2^{(1)}(q^{(2)})$  given by (5.13) and (5.14), such that

$$Q^{(1)}(k;m) \cap (r_2 + J_{k+16}(\ell_3)) = \emptyset.$$
(5.17)

Since  $J_{k+16}(\ell_3) \subset J_{k+8}(q^{(2)})$ , it clearly follows from (5.15) and (5.16) that

$$Q^{(1)}(k;m) \cap (r_2^{(0)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \tag{5.18}$$

$$Q^{(1)}(k;m) \cap (r_2^{(1)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset.$$
 (5.19)

Since (5.17)–(5.19) are analogs of (5.4), the *T*-power expansion argument shows that for every integer  $1 - \ell_3 \leq h \leq q_{k+17} - 2 - \ell_3$ , we have

$$Q^{(2)}(k;m) \cap T^h(r_2 + J_{k+16}(\ell_3)) = \emptyset, \tag{5.20}$$

$$Q^{(2)}(k;m) \cap T^{h}(r_2^{(0)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \tag{5.21}$$

$$Q^{(2)}(k;m) \cap T^h(r_2^{(1)}(q^{(2)}) + J_{k+16}(\ell_3)) = \emptyset, \tag{5.22}$$

where, corresponding to  $Q^{**}(k; m)$  in Section 4, we write

$$Q^{(2)}(k;m) = \{ y_i : q_{k+1} + q_{k+9} + q_{k+17} \leqslant i \leqslant m - q_{k+1} - q_{k+9} - q_{k+17} \}.$$

For notational convenience, for every integer  $1 - \ell_3 \leqslant h \leqslant q_{k+17} - 2 - \ell_3$ , we write

$$T^{h}(r_{2}^{(0)}(q^{(2)}) + J_{k+16}(\ell_{3})) = r_{3}^{(0)}(\ell_{3} + h) + J_{k+16}(\ell_{3} + h), \tag{5.23}$$

$$T^{h}(r_{2}^{(1)}(q^{(2)}) + J_{k+16}(\ell_{3})) = r_{3}^{(1)}(\ell_{3} + h) + J_{k+16}(\ell_{3} + h), \tag{5.24}$$

$$T^{h}(r_{2} + J_{k+16}(\ell_{3})) = r_{3}^{(2)}(\ell_{3} + h) + J_{k+16}(\ell_{3} + h).$$
 (5.25)

Then combining (5.20)–(5.25), we have

$$Q^{(2)}(k;m) \cap (r_3^{(0)}(q) + J_{k+16}(q) = \emptyset,$$

$$Q^{(2)}(k;m) \cap (r_3^{(1)}(q) + J_{k+16}(q) = \emptyset,$$

$$Q^{(2)}(k;m) \cap (r_3^{(2)}(q) + J_{k+16}(q) = \emptyset,$$

for every integer  $1 \leq q \leq q_{k+17} - 2$ . Clearly

$$r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q)$$
 are distinct,  $1 \le q \le q_{k+17} - 2$ .

We now split Case 2B into two complementary cases.

Case 3A. The following intersection property holds. For every

$$J_{k+24}(\ell) \subset J_{k+16}(q)$$
 and  $r \in \{0, 1, \dots, s-1\},\$ 

with  $r \neq r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q)$  given by (5.23)–(5.25), we have

$$Q^{(2)}(k;m) \cap (r + J_{k+24}(\ell)) \neq \emptyset.$$

Case 3B. There exist

$$J_{k+24}(\ell_4) \subset J_{k+16}(q^{(3)})$$
 and  $r_3 \in \{0, 1, \dots, s-1\},$ 

with  $r_3 \neq r_3^{(0)}(q), r_3^{(1)}(q), r_3^{(2)}(q)$  given by (5.23)–(5.25), such that

$$Q^{(2)}(k;m) \cap (r_3 + J_{k+24}(\ell_4)) = \emptyset.$$

Suppose that  $1 \le \tau \le s - 2$ . Assume that for every integer  $1 \le q \le q_{k+8\tau-7} - 2$ , there are distinct integers

$$r_{\tau}^{(0)}(q), \dots, r_{\tau}^{(\tau-1)}(q) \in \{0, 1, \dots, s-1\}$$
 (5.26)

such that for every integer  $1 \leq j \leq \tau$ ,

$$Q^{(\tau-1)}(k;m) \cap (r_{\tau}^{(j-1)}(q) + J_{k+8\tau-8}(q)) = \emptyset$$
 (5.27)

for every integer  $1 \leqslant q \leqslant q_{k+8\tau-7} - 2$ , where

$$Q^{(\tau-1)}(k;m) = \left\{ y_i : \sum_{u=1}^{\tau} q_{k+8u-7} \leqslant i \leqslant m - \sum_{u=1}^{\tau} q_{k+8u-7} \right\}.$$

Assume further that we have two complementary cases.

Case  $\tau A$ . The following intersection property holds. For every

$$J_{k+8\tau}(\ell) \subset J_{k+8\tau-8}(q)$$
 and  $r \in \{0, 1, \dots, s-1\},$ 

with  $r \neq r_{\tau}^{(0)}(q), \dots, r_{\tau}^{(\tau-1)}(q)$  given by (5.26), we have

$$Q^{(\tau-1)}(k;m) \cap (r + J_{k+8\tau}(\ell)) \neq \emptyset.$$

Case  $\tau B$ . There exist

$$J_{k+8\tau}(\ell_{\tau+1}) \subset J_{k+8\tau-8}(q^{(\tau)})$$
 and  $r_{\tau} \in \{0, 1, \dots, s-1\},$ 

with  $r_{\tau} \neq r_{\tau}^{(0)}(q), \dots, r_{\tau}^{(\tau-1)}(q)$  given by (5.26), such that

$$Q^{(\tau-1)}(k;m) \cap (r_{\tau} + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset.$$
 (5.28)

Since  $J_{k+8\tau}(\ell_{\tau+1}) \subset J_{k+8\tau-8}(q^{(\tau)})$ , it clearly follows from (5.27) that for every integer  $1 \leq j \leq \tau$ ,

$$Q^{(\tau-1)}(k;m) \cap (r_{\tau}^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset.$$
 (5.29)

Since (5.28) and (5.29) are analogs of (5.4), the *T*-power expansion argument shows that for every integer  $1 - \ell_{\tau+1} \leq h \leq q_{k+8\tau+1} - 2 - \ell_{\tau+1}$ , we have

$$Q^{(\tau)}(k;m) \cap T^h(r_{\tau} + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset, \tag{5.30}$$

as well as

$$Q^{(\tau)}(k;m) \cap T^h(r_{\tau}^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = \emptyset$$
(5.31)

for every integer  $1 \leq j \leq \tau$ , where

$$Q^{(\tau)}(k;m) = \left\{ y_i : \sum_{u=1}^{\tau+1} q_{k+8u-7} \leqslant i \leqslant m - \sum_{u=1}^{\tau+1} q_{k+8u-7} \right\}.$$

For notational convenience, for every integer  $1 - \ell_{\tau+1} \leq h \leq q_{k+8\tau+1} - 2 - \ell_{\tau+1}$ , we write

$$T^{h}(r_{\tau}^{(j-1)}(q^{(\tau)}) + J_{k+8\tau}(\ell_{\tau+1})) = r_{\tau+1}^{(j-1)}(\ell_{\tau+1} + h) + J_{k+8\tau}(\ell_{\tau+1} + h)$$
 (5.32)

for every integer  $1 \leq j \leq \tau$ , and also write

$$T^{h}(r_{\tau} + J_{k+8\tau}(\ell_{\tau+1})) = r_{\tau+1}^{(\tau)}(\ell_{\tau+1} + h) + J_{k+8\tau}(\ell_{\tau+1} + h). \tag{5.33}$$

Then combining (5.30)–(5.33), we have, for every integer  $1 \le j \le \tau + 1$ ,

$$Q^{(\tau)}(k;m) \cap (r_{\tau+1}^{(j-1)}(q) + J_{k+8\tau}(q)) = \emptyset$$

for every integer  $1 \leq q \leq q_{k+8\tau+1} - 2$ . Clearly

$$r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q)$$
 are distinct,  $1 \leqslant q \leqslant q_{k+8\tau+1} - 2$ .

We now split Case  $\tau B$  into two complementary cases.

Case  $(\tau + 1)A$ . The following intersection property holds. For every

$$J_{k+8\tau+8}(\ell) \subset J_{k+8\tau}(q)$$
 and  $r \in \{0, 1, \dots, s-1\},\$ 

with  $r \neq r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q)$  given by (5.32) and (5.33), we have

$$Q^{(\tau)}(k;m) \cap (r + J_{k+8\tau+8}(\ell)) \neq \emptyset.$$

Case  $(\tau + 1)B$ . There exist

$$J_{k+8\tau+8}(\ell_{\tau+2}) \subset J_{k+8\tau}(q^{(\tau+1)})$$
 and  $r_{\tau+1} \in \{0, 1, \dots, s-1\},$ 

with  $r_{\tau+1} \neq r_{\tau+1}^{(0)}(q), \dots, r_{\tau+1}^{(\tau)}(q)$  given by (5.32) and (5.33), such that

$$Q^{(\tau)}(k;m) \cap (r_{\tau+1} + J_{k+8\tau+8}(\ell_{\tau+2})) = \emptyset.$$

In particular, if  $\tau = s - 2$ , we have the following final case.

Case (s-1)B. There exist

$$J_{k+8s-8}(\ell_s) \subset J_{k+8s-16}(q^{(s-1)})$$
 and  $r_{s-1} \in \{0, 1, \dots, s-1\},$ 

with  $r_{s-1} \neq r_{s-1}^{(0)}(q^{(s-1)}), \ldots, r_{s-1}^{(s-2)}(q^{(s-1)})$  given by (5.32) and (5.33) in the special case  $\tau = s - 2$ , such that

$$Q^{(s-2)}(k;m) \cap (r_{s-1} + J_{k+8s-8}(\ell_s)) = \emptyset.$$
 (5.34)

**Lemma 5.2.** For every  $\tau = 1, ..., s - 1$ , Case  $\tau A$  is impossible.

**Lemma 5.3.** If Case (s-1)B holds, then

$$m \le 2s + 2\sum_{u=1}^{s} q_{k+8u-7}.$$
 (5.35)

Before we prove Lemmas 5.2 and 5.3, we first complete the proof of Theorem 1.

Proof of Theorem 1. Suppose that  $I_0 \subset [0,s)$  is  $(\mathcal{L}_{\alpha}; M)$ -free, so that

$$\{y_i: 1 \leqslant i \leqslant m\} \cap I_0 = \emptyset.$$

Let  $J_k(\ell_1) = J(\alpha; k; \ell_1)$  be the longest special interval of the form (3.5) such that

$$r(\ell_1) + J_k(\ell_1) \subset I_0$$
 for some  $\ell_1$  and  $r(\ell_1) \in \{0, 1, \dots, s-1\}$ .

Then we can show as before that

$$\operatorname{length}(I_0) < \frac{8}{q_k}. \tag{5.36}$$

On the other hand, it follows from (5.1) and (5.2) that

$$M \leqslant (m+1)\sqrt{1+\alpha^2}.\tag{5.37}$$

Also, in view of Lemmas 5.2 and 5.3, it is clear that the bound (5.35) holds. Finally, the inequalities (3.33) are replaced by the inequalities

$$q_{k+1} < q_{k+9} < \dots < q_{k+8s-7} \le (A+1)^{8s-7} q_k.$$
 (5.38)

Combining (5.35), (5.37) and (5.38), we see that

$$M \leqslant \left(2s + 1 + 2\sum_{u=1}^{s} q_{k+8u-7}\right)\sqrt{1+\alpha^2} < (4s+1)(A+1)^{8s-7}q_k\sqrt{2}.$$
 (5.39)

It now follows from (5.36) and (5.39) that a geodesic segment  $\mathcal{L}_{\alpha}(t)$ ,  $0 \leq t \leq M$ , of length  $M = (4s+1)(A+1)^{8s-7}q_k\sqrt{2}$  must intersect every subinterval I of  $h_1 \cup \ldots \cup h_s$  with length I =  $8/q_k$ . Since the product I length I is a constant independent of I, this establishes superdensity of the half-infinite geodesic.

It remains to prove Lemmas 5.2 and 5.3.

Proof of Lemma 5.2. We can proceed along similar lines as in the first part of the proof of Lemma 3.2, and show that any two neighboring s-copy extensions  $J_{k+8\tau-8}(q';s)$  and  $J_{k+8\tau-8}(q'';s)$  are synchronized in the following precise sense: For each  $r \in \{0,1,\ldots,s-1\}$ , we have

$$Q^{(\tau-1)}(k;m) \cap (r + J_{k+8\tau-8}(q')) = \emptyset$$

if and only if

$$Q^{(\tau-1)}(k;m) \cap (r + J_{k+8\tau-8}(q'')) = \emptyset.$$

By definition, the s-copy extensions  $J_{k+8\tau-8}(q;s)$ ,  $1 \leq q \leq q_{k+8\tau-7} - 2$ , give rise to 2s continuous chains of overlapping intervals in the torus/circle [0,s) such that the 2s chains completely cover the 2s intervals

$$[0, 1 - \alpha), [1 - \alpha, 1), \dots, [s - 1, s - \alpha), [s - \alpha, s).$$
 (5.40)

The synchronization property now implies that each of the 2s long special intervals in (5.40) satisfies one of the following two properties. Either such a long special interval is disjoint from the set  $Q^{(\tau-1)}(k;m)$ , or the set  $Q^{(\tau-1)}(k;m)$  is dense in such a long special interval, in the precise sense that every subinterval of length  $1/q_{k+8s}$  contains a point from the set  $Q^{(\tau-1)}(k;m)$ .

Moreover, it is not difficult to show that precisely  $2\tau$  of the 2s long special intervals in (5.40) are disjoint from the set  $Q^{(\tau-1)}(k;m)$ . To see this, choose two integers q' and q'' satisfying  $1 \leq q', q'' \leq q_{k+8\tau-7} - 2$  such that

$$J_{k+8\tau-8}(q') \subset (0, 1-\alpha)$$
 and  $J_{k+8\tau-8}(q'') \subset (1-\alpha, 1)$ . (5.41)

Then it follows from (5.27) that for every integer  $1 \leq j \leq \tau$ ,

$$Q^{(\tau-1)}(k;m) \cap (r_{\tau}^{(j-1)}(q') + J_{k+8\tau-8}(q')) = \emptyset, \tag{5.42}$$

$$Q^{(\tau-1)}(k;m) \cap (r_{\tau}^{(j-1)}(q'') + J_{k+8\tau-8}(q'')) = \emptyset.$$
 (5.43)

For every integer  $1 \leq j \leq \tau$ , now write

$$\mathcal{I}_{2j-1} = r_{\tau}^{(j-1)}(q') + (0, 1 - \alpha) \quad \text{and} \quad \mathcal{I}_{2j} = r_{\tau}^{(j-1)}(q'') + (1 - \alpha, 1).$$
 (5.44)

The synchronization property and (5.41)–(5.44) now imply that for every integer  $1 \le j \le \tau$ , we have

$$Q^{(\tau-1)}(k;m) \cap \mathcal{I}_{2j-1} = \emptyset$$
 and  $Q^{(\tau-1)}(k;m) \cap \mathcal{I}_{2j} = \emptyset$ .

Note that the union  $\mathcal{I}_1 \cup \ldots \cup \mathcal{I}_{2\tau}$  modulo 1 is precisely the unit interval [0,1) taken  $\tau$  times.

Now  $\mathcal{I}_1, \ldots, \mathcal{I}_{2\tau}$  are  $2\tau$  of the 2s long special intervals in (5.40). Let the remaining long special intervals in (5.40) be denoted by  $\mathcal{I}_{2j-1}$  and  $\mathcal{I}_{2j}$ ,  $\tau < j \leq s$ . The condition of Case  $\tau A$  now implies that these  $2s-2\tau$  intervals are not disjoint from  $Q^{(\tau-1)}(k;m)$ , so that  $Q^{(\tau-1)}(k;m)$  is dense in each of them.

Each T-image  $T(\mathcal{I}_{2j-1})$  and  $T(\mathcal{I}_{2j})$ ,  $1 \leq j \leq \tau$ , has at most 1 common point with the set  $Q^{(\tau-1)}(k;m)$ . This is a contradiction, since the union

$$T(\mathcal{I}_1) \cup \ldots \cup T(\mathcal{I}_{2\tau})$$

has a *substantial* intersection with the union  $\mathcal{I}_{2\tau+1} \cup \ldots \cup \mathcal{I}_{2s}$ , which implies that it must have a *substantial* intersection with the set  $Q^{(\tau-1)}(k;m)$ , much more than at most  $2\tau$  elements. Thus Case  $\tau A$  is impossible, and this completes the proof.

Proof of Lemma 5.3. Since (5.34) is an analog of (5.4), we can repeat the *T*-power extension argument and conclude that for every integer  $1 - \ell_s \leq h \leq q_{k+8s-7} - 2 - \ell_s$ , we have

$$Q^{(s-1)}(k;m) \cap T^h(r_{s-1} + J_{k+8s-8}(\ell_s)) = \emptyset.$$
 (5.45)

Since  $J_{k+8s-8}(\ell_s) \subset J_{k+8s-16}(q^{(s-1)})$ , it follows from (5.27) with  $\tau = s-1$  that for every integer  $1 \leq j \leq s-1$ ,

$$Q^{(s-2)}(k;m) \cap (r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = \emptyset.$$
 (5.46)

Next, note that (5.46) are also analogs of (5.4), so again we can repeat the T-power extension argument and conclude that for every integer  $1 \le j \le s-1$ , and for every integer  $1 - \ell_s \le h \le q_{k+8s-7} - 2 - \ell_s$ , we have

$$Q^{(s-1)}(k;m) \cap T^h(r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = \emptyset.$$
 (5.47)

Now, for every integer  $1 - \ell_s \leq h \leq q_{k+8s-7} - 2 - \ell_s$ , we write

$$T^{h}(r_{s-1}^{(j-1)}(q^{(s-1)}) + J_{k+8s-8}(\ell_s)) = r^{(j-1)}(\ell_s + h) + J_{k+8s-8}(\ell_s + h)$$
(5.48)

for every integer  $1 \leq j \leq s-1$ , and also write

$$T^{h}(r_{s-1} + J_{k+8s-8}(\ell_s)) = r^{(s-1)}(\ell_s + h) + J_{k+8s-8}(\ell_s + h).$$
 (5.49)

Clearly it follows from the assumption of Case (s-1)B that  $r^{(0)}(q), \ldots, r^{(s-1)}(q)$  form a permutation of  $0, 1, \ldots, s-1$  for every integer  $1 \leq q \leq q_{k+8s-7}-2$ . Combining (5.45) and (5.47)–(5.49), we have, for every integer  $1 \leq j \leq s$ ,

$$Q^{(P-1)}(k;m) \cap (r^{(j-1)}(q) + J_{k+8s-8}(q)) = \emptyset$$
 (5.50)

for every integer  $1 \leqslant q \leqslant q_{k+8s-7} - 2$ .

Note now that (5.50) are similar to (4.7)–(4.9) in the proof of Lemma 4.1, so we now mimic the last part of that proof.

By definition, the s-copy extensions  $J_{k+8s-8}(q;s)$ ,  $1 \le q \le q_{k+8s-7} - 2$ , give rise to 2s continuous chains of overlapping intervals in the torus/circle [0,s) such that the 2s chains completely cover the 2s intervals

$$[0, 1 - \alpha), [1 - \alpha, 1), \ldots, [s - 1, s - \alpha), [s - \alpha, s),$$

and there are only 2s points in [0, s) that are not covered by the 2s chains, namely

$$j-1$$
,  $j-\alpha$ ,  $1 \le j \le s$ .

Combining (5.50) for every integer  $1 \leq q \leq q_{k+8s-7} - 2$ , we deduce that the set  $Q^{(s-1)}(k;m)$  is not covered by the 2s chains. Indeed, if

$$m \geqslant 2s + 1 + 2\sum_{u=1}^{s} q_{k+8u-7},$$

then the set  $Q^{(s-1)}(k;m)$  has at least 2s+1 distinct elements, which is more than 2s, giving rise to a contradiction. We conclude therefore that, under the conditions of Case (s-1)B, we must have

$$m \leqslant 2s + 2\sum_{u=1}^{s} q_{k+8u-7},$$

and this completes the proof.

### 6. A DYNAMICAL SYSTEMS APPROACH

Consider general translation surfaces, *i.e.*, polygons or finite sets of polygons on the plane such that each side of a polygon is identified with a parallel side by translation. There is a long history of links between the dynamical systems on individual translation surfaces and a dynamical system on the moduli space of translation surfaces, and between the linear flow on a translation surface and geodesic flow on the moduli space. Central to this is Masur's criterion on unique ergodicity which is built on his earlier work with Kerckhoff and Smillie; see [7, 10, 11].

Since the appearance of an earlier version of this paper on arXiv, Southerland [15] has studied the problem involving general translation surfaces and established a

beautiful geometric result that a linear flow on a translation surface is superdense if and only if the associated Teichmüller geodesic trajectory is bounded in the moduli space. This raises the question of the relationship between the boundedness of the associated Teichmüller geodesic trajectory in the moduli space and the arithmetic properties of the slope of the linear flow on the translation surface.

This relationship is unclear other than what is evidenced on finite polysquare translation surfaces and what happens in the moduli space of tori. Does the set of slopes that lead to superdense linear flow on a translation surface change if the surface looks less like a finite polysquare translation surface? Is the torus, and by extension, finite polysquare translation surfaces maximal in some sense?

Acknowledgments. The authors wish to thank the referee for the very careful reading of the manuscript and for the many valuable comments.

### REFERENCES

- [1] J. Beck, M. Donders, Y. Yang. Quantitative behavior of non-integrable systems (I). *Acta Math. Hungar.* **161** (2020), 66–184.
- [2] J. Beck, M. Donders, Y. Yang. Quantitative behavior of non-integrable systems (II). Acta Math. Hungar. 162 (2020), 220–324.
- [3] J. Beck, W.W.L. Chen, Y. Yang. Quantitative behavior of non-integrable systems (III). *Acta Math. Hungar.* **166** (2022), 254–372.
- [4] J. Beck, W.W.L. Chen, Y. Yang. Quantitative behavior of non-integrable systems (IV). *Acta Math. Hungar.* **167** (2022), 1–160.
- [5] J.H. Halton. The distribution of the sequence  $\{n\xi\}$  (n = 0, 1, 2, ...). Proc. Cambridge Philos. Soc. **61** (1965), 665–670.
- [6] A. Katok, A. Zemlyakov. Topological transitivity of billiards in polygons. *Math. Notes* **18** (1975), 760–764.
- [7] S. Kerckhoff, H. Masur, J. Smillie. Ergodicity of billiard flows and quadratic differentials. *Ann. of Math.* **124** (1986), 293–311.
- [8] A.Ya. Khinchin. Continued Fractions (Dover, 1997).
- [9] D. König, A. Szücs. Mouvement d'un point abondonne a l'interieur d'un cube. *Rend. Circ. Mat. Palermo* **36** (1913), 79–90.
- [10] H. Masur. Hausdorff dimension of the set of nonergodic foliations of a quadratic differential. Duke Math. J. 66 (1992), 387–442.
- [11] H. Masur, J. Smillie. Hausdorff dimension of sets of nonergodic measured foliations. *Ann. of Math.* **134** (1991), 455–543.
- [12] N.B. Slater. Gap and steps for the sequence  $n\theta \mod 1$ . Proc. Cambridge Philos. Soc. **63** (1967), 1115-1123.
- [13] V.T. Sós. On the theory of diophantine approximations. *Acta Math. Acad. Sci. Hungar.* 8 (1957), 461–472.
- [14] V.T. Sós. On the distribution mod 1 of the sequence  $n\alpha$ . Ann. Univ. Sci. Budapest Eötvös Sect. Math. 1 (1958), 127–134.
- [15] J. Southerland. Superdensity and bounded geodesic in moduli space (preprint). arxiv.org/abs/2201.10156
- [16] J. Surányi. Über die Anordnung der Vielfachen einer reellen Zahl mod 1. Ann. Univ. Sci. Budapest Eötvös Sect. Math. 1 (1958), 107–111.
- [17] S. Swierczkowski. On successive settings of an arc on the circumference of a circle. *Fund. Math.* **46** (1959), 187–189.

DEPARTMENT OF MATHEMATICS, HILL CENTER FOR THE MATHEMATICAL SCIENCES, RUTGERS UNIVERSITY, PISCATAWAY NJ 08854, USA

 $E ext{-}mail\ address: jbeck@math.rutgers.edu}$ 

School of Mathematical and Physical Sciences, Faculty of Science and Engineering, Macquarie University, Sydney NSW 2109, Australia

E-mail address: william.chen@mq.edu.au